

COVID-19 vaccines for people with a weakened immune system



North Central London
Clinical Commissioning Group

If you are....	You can get....	What are the criteria?
Aged 5-11 with a health condition that puts you at higher risk from COVID-19, or you live with someone with a weakened immune system	<ol style="list-style-type: none"> 1. First dose and second dose, at least 8 weeks apart 	Conditions and treatments are listed in Table 4 on page 22 of the Green Book . All of Table 4 applies for this category.
Aged 12-15 with a health condition that puts you at higher risk from COVID-19, or you live with someone who has a weakened immune system	<ol style="list-style-type: none"> 1. First dose and second dose, at least eight weeks apart 2. Booster dose from three months (91 days) after previous dose 	Conditions and treatments are listed in Table 4 on page 22 of the Green Book . All of Table 4 applies for this category.
Aged 12+ and have a weakened immune system	<ol style="list-style-type: none"> 1. First dose and second dose, at least eight weeks apart 2. Booster dose from three months (91 days) after previous dose 3. Spring booster around 6 months (and not before 3 months) after previous dose. 	<p>For ages 12-15, conditions and treatments are listed in the “immunosuppression” row of Table 4 on page 22 of the Green Book</p> <p>For ages 16+, conditions and treatments are listed in the “immunosuppression” row of Table 3 on page 16 of the Green Book</p>
Aged 12+ and have a severely weakened immune system	<ol style="list-style-type: none"> 1. First dose and second dose, at least eight weeks apart 2. Third primary dose, typically eight weeks after second dose (the timing may vary depending on any treatment you may be having) 3. Booster dose from three months (91 days) after previous dose 4. Spring booster around 6 months (and not before 3 months) after previous dose. 	Conditions and treatments are listed in Box 1 on page 25 of Green Book .